

Black Darter

Sympetrum danae

Insects · On the wing Jul-Oct



Photo by S. Rae from Scotland, UK, CC BY 2.0, via Wikimedia Commons

A small dragonfly. Mature males turn almost entirely matt black with a few yellow flecks; females and immatures are yellow and black with a distinctive black triangle on top of the thorax. Black darters are birds of the bog — they need acidic, peaty pools, mires and moorland lochans, which makes the peatlands of Dumfries and Galloway good ground for them.

BEHAVIOUR

Like other darters it hunts from a low perch, often on bare peat, sphagnum or heather, returning repeatedly to the same spot. It is strongly tied to acid water with abundant sphagnum moss, so it is a useful indicator that a bog is in reasonable condition. Larvae develop in a single year in shallow, warm, peaty pools that dry out too often for larger species.

WHEN TO SEE IT

J	F	M	A	M	J	J	A	S	O	N	D
---	---	---	---	---	---	---	---	---	---	---	---

QUICK FACTS

FAMILY	Skimmers and darters (Libellulidae)
SIZE	Body 29-34 mm · Britain's smallest dragonfly
STATUS	Native; a peatland specialist, commonest in Scotland

Photo by S. Rae from Scotland, UK, CC BY 2.0, via Wikimedia Commons

Natureville