

Common Blue

Polyommatus icarus

Insects · On the wing May-Sep · two broods



Photo by Charles J. Sharp, CC BY-SA 4.0, via Wikimedia Commons

Our commonest blue butterfly. Males are a bright violet-blue above with a fine white fringe; females are brown, usually with a dusting of blue at the wing bases and a row of orange crescents along the margins, and are frequently mistaken for a different species. Both sexes share a grey-brown underside spotted with black and orange. Found on unimproved grassland, verges and dunes around Dumfries.

BEHAVIOUR

Common blues depend entirely on bird's-foot trefoil and related vetches, so they disappear from over-fertilised or heavily grazed grassland. Adults roost communally, several individuals settling head-down on the same grass stem overnight. The caterpillars secrete a sugary fluid and are attended and protected by ants, a relationship shared with many of the blues.

WHEN TO SEE IT

J	F	M	A	M	J	J	A	S	O	N	D
---	---	---	---	---	---	---	---	---	---	---	---

QUICK FACTS

FAMILY	Blues and coppers (Lycaenidae)
WINGSPAN	3–3.5 cm
STATUS	Native; widespread on unimproved grassland

Photo by Charles J. Sharp, CC BY-SA 4.0, via Wikimedia Commons
Natureville